

WEST Search History

DATE: Tuesday, November 12, 2002

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=USPT; PLUR=YES; OP=AND</i>			
L19	L18 and galectin	0	L19
L18	l16 and L17	49	L18
L17	daniel.in.	36166	L17
L16	hsu.in.	3716	L16
L15	L14 and galectin	0	L15
L14	l12 and L13	37	L14
L13	hideki.in.	5973	L13
L12	sano.in.	2371	L12
L11	l9 and L10	2	L11
L10	fu-tong.in.	4	L10
L9	liu.in.	5535	L9
L8	monocyte or macrophage	18454	L8
L7	l3 and L6	17	L7
L6	migration	56872	L6
L5	l3 and L4	8	L5
L4	cell with migration	6367	L4
L3	l1 or L2	27	L3
L2	galectin adj 3	27	L2
L1	galectin-3	26	L1

END OF SEARCH HISTORY

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 11:38:35 ON 12 NOV 2002

=> file caplus medline biosis

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'CAPLUS' ENTERED AT 11:38:44 ON 12 NOV 2002
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'MEDLINE' ENTERED AT 11:38:44 ON 12 NOV 2002

FILE 'BIOSIS' ENTERED AT 11:38:44 ON 12 NOV 2002
COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC. (R)

=> galectin 3

L1 1152 GALECTIN 3

=> cell(s)migrat?

L2 120256 CELL(S) MIGRAT?

=> 11 and 12

L3 61 L1 AND L2

=> 13 and 1970-2000/py

L4 40 L3 AND 1970-2000/PY

=> monocyte or macrophage

L5 447251 MONOCYTE OR MACROPHAGE

=> 11 and 15

L6 160 L1 AND L5

=> 13 and 15

L7 11 L3 AND L5

=> 14 and 15

L8 8 L4 AND L5

=> dup rem l8

PROCESSING COMPLETED FOR L8
L9 4 DUP REM L8 (4 DUPLICATES REMOVED)

=> d ti abs so 19 1-4

L9 ANSWER 1 OF 4 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 1
 TI Human **galectin-3** is a novel chemoattractant for
monocytes and **macrophages**
 AB **Galectin-3** is a .beta.-galactoside-binding protein
 implicated in diverse biol. processes. The authors found that
galectin-3 induced human **monocyte** migration in
 vitro in a dose-dependent manner, and it was chemotactic at high concns.
 (1.0 .mu.M) but chemokinetic at low concns. (10-100 nM). **Galectin**
-3-induced **monocyte** migration was inhibited by its
 specific mAb and was blocked by lactose and a C-terminal domain fragment
 of the protein, indicating that both the N-terminal and C-terminal
 domains
 of **galectin-3** are involved in this activity.
 Pertussis toxin (PTX) almost completely blocked **monocyte**
 migration induced by high concns. of **galectin-3**.
Galectin-3 caused a Ca²⁺ influx in **monocytes**
 at high, but not low, concns., and both lactose and PTX inhibited this
 response. There was no cross-desensitization between **galectin-**
3 and any of the **monocyte**-reactive chemokines examd.,
 including **monocyte** chemotactic protein-1, **macrophage**
 inflammatory protein-1.alpha., and stromal cell-derived factor-1.alpha..
 Cultured human **macrophages** and alveolar **macrophages**
 also migrated toward **galectin-3**, but not
monocyte chemotactic protein-1. Finally, **galectin-**
3 was found to cause **monocyte** accumulation in vivo in
 mouse air pouches. Thus, **galectin-3** is a novel
 chemoattractant for **monocytes** and **macrophages** and its
 effect is mediated at least in part through a PTX-sensitive (G
 protein-coupled) pathway.
 SO Journal of Immunology (2000), 165(4), 2156-2164
 CODEN: JOIMA3; ISSN: 0022-1767

L9 ANSWER 2 OF 4 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 2
 TI **Galectin-3** gene (LGALS3) expression in experimental
 atherosclerosis and cultured smooth muscle cells
 AB The **galectin-3** gene (LGALS3) encodes a
 .beta.-galactose-binding lectin. LGALS3 expression is assocd. with
 neoplastic transformation and with differentiation of **monocytes**
 to **macrophages**. Factors involved in **migration**,
 proliferation, adhesion, and differentiation of vascular smooth muscle
cells (SMC) play a major role during atherosclerosis development.
 Expression of the **galectin-3** gene was not detected in
 quiescent SMC but was activated in aortas of hypercholesterolemic
 rabbits,
 in aortas of rats after balloon injury, and in cultured SMC. Thus,
galectin-3 prodn. is involved in the developmental
 process of atherogenesis.
 SO FEBS Letters (1998), 430(3), 307-311
 CODEN: FEBLAL; ISSN: 0014-5793

L9 ANSWER 3 OF 4 CAPLUS COPYRIGHT 2002 ACS
 TI Maintenance of granulocyte numbers during acute peritonitis is defective
 in **galectin-3**-null mutant mice
 AB **Galectin-3**, also known as the **macrophage**
 marker Mac-2, is a member of a family of structurally related animal
 lectins that exhibit specificity for .beta.-galactosides. To investigate
 the role of **galectin-3** in acute inflammation, the
 authors compared the no. of leukocytes present in the peritoneal cavity
 of
 wild type and **galectin-3** null mutant mice after i.p.

injection of thioglycolate broth. At day 1 after injection, the authors found no difference in the recruitment of mononuclear phagocytes and granulocytes to the peritoneal cavity. However, 4 days after thioglycolate injection, **galectin-3** mutant mice exhibited a significantly reduced no. of recoverable granulocytes compared to wild-type animals. As mutant granulocytes did not exhibit an accelerated rate of apoptosis and their uptake by **macrophages** appeared to be unaffected by the mutation, the phenotype described here suggests that **galectin-3** participates in an addnl. level of control during the resoln. of acute inflammation.

SO Immunology (1998), 94(3), 290-296
CODEN: IMMUAM; ISSN: 0019-2805

L9 ANSWER 4 OF 4 MEDLINE DUPLICATE 3

TI Soybean agglutinin binds a 160-kDa rat **macrophage** membrane glycoprotein and enhances cell differentiation and activation.

AB Mature **macrophages** (M phi) differ from other rat leukocytes by their ability to bind soybean agglutinin (SBA). In this study we identify the SBA-binding structure on rat bone marrow-derived M phi (BMDM phi). Precipitation of iodinated membrane proteins from rat bone marrow **cells** (BMC) and BMDM phi with SBA revealed a major glycoprotein of Mr 160 kDa on BMDM phi but not on BMC. In addition minor bands **migrating** at 70 and 26 kDa were seen. Stimulation of BMDM phi with 100 nM SBA induced a decrease in surface density of Thy1.1 (MRC OX7) and His54 and an increase in the expression of MRC OX6 (RT1.B/I-A), MRC OX17 (RT1.D/I-E), MRC OX41 (gp 110/120), MRC OX42 (CD11b/c), Mac1 (CD11b/CR3) and Mac2 (**galectin-3**/IgE binding protein) antigen. Expression of other M phi differentiation antigens recognized by mAb MRC OX43 (M phi, endothelial **cells**) and ED9 (M phi/CD14 like) were not significantly altered. BMDM phi derived from cultures with M phi colony-stimulating factor (M-CSF) and SBA showed increased oxidative burst and phagocytic activity compared to **cells** cultured with M-CSF alone. Our data suggest that binding of a 160-kDa membrane glycoprotein on M phi by N-acetylgalactosamine-specific lectins stimulates M phi differentiation and activation.

SO IMMUNOLOGY LETTERS, (1996 Aug) 52 (1) 53-6.
Journal code: 7910006. ISSN: 0165-2478.

=> logoff

FAOM Checklist

Appl. Ser. No.

09/805449

FD

3/13/01

5

date

EFD

3/13/00

via

60/188795



Briefed

Form 948 present?

10



Declaration (06-05)

PCT referred to?

priority dates OK?

continuing dates OK?

15

CIP duty? (06-05-09)



Meets conditions for priority (02-09)

W/A

Sequences in compliance?

20



Restriction/Election?

NA

Priority Docs present?

25



File wrapper labeled OK and initialed?

obj

Title (06-11)



Abstract (06-12 missing)

(06-13 informs)

(06-14, 15 content)

30



Figures (06-27)

35



Brief Description



Amendments?

na

IDS checked

40

na

Seq Srch requested



Copies of refs requested

45



07-29, 06-31?



07-34